

Quadratic Functions and Modeling

In this unit, we will study quadratic functions and the relationships for which they provide suitable models. An important application of such functions is to describe the trajectory, or path, of an object near the surface of the earth when the only force acting on the object is gravitational attraction. What happens when you toss a ball straight up into the air? What about an outfielder on a baseball team throwing a ball into the infield? If air resistance and outside forces are negligible, what is the mathematical model for the relationship between time and height of the ball?

Definition: *Quadratic Functions*

A **quadratic function** is one of the form

$$f(x) = ax^2 + bx + c,$$

where a , b , and c are real numbers with $a \neq 0$.

The graph of a quadratic function is called a ***parabola*** and its shape resembles that of the graph in each of the following two examples.

Example 1

Figure 1 shows the graph of the quadratic function

$$y = f(x) = x^2 - 4x + 1.$$




figure 1

Observe that there is a lowest point $V(2, -3)$ on the graph in figure 1. The point V is called the ***vertex*** of the parabola.

2
Quadratic Functions and Modeling

Example 2

Figure 2 shows the graph of the quadratic function

$$y = g(x) = -2x^2 + 4x + 3.$$




figure 2

Again, observe that there is a highest point $V(1, 5)$ on the graph in figure 2. This point V is also the ***vertex*** of the parabola.

By **completing the square**,

the quadratic function (example 1) $f(x) = x^2 - 4x + 1$ can be written as

$$f(x) = (x - 2)^2 - 3;$$

the quadratic function (example 2) $g(x) = -2x^2 + 4x + 3$ can be written as

$$g(x) = -2(x - 1)^2 + 5.$$

Note that the parabola in example 1 opens upward, with vertex $V(2, -3)$ and a vertical axis of symmetry $x = 2$. The parabola in example 2 opens downward with vertex $V(1, 5)$ and a vertical axis of symmetry $x = 1$.

Quadratic Functions and Modeling

Standard Form of a Quadratic Function Whose Graph (a Parabola) has a Vertical Axis

The graph of $f(x) = ax^2 + bx + c$, $a \neq 0$
 $= a(x - h)^2 + k$

is a parabola that has vertex $V(h, k)$ and a vertical axis.

The parabola opens upward if $a > 0$ or downward if $a < 0$.

If $a > 0$, the vertex $V(h, k)$ is the lowest point on the parabola, and the function f has a minimum value $f(h) = k$.

If $a < 0$, the vertex $V(h, k)$ is the highest point on the parabola, and the function f has a maximum value $f(h) = k$.

Quadratic Models and Equations

Quadratic Population Models:

$$P(t) = P_0 + bt + at^2.$$

Here, we denote the independent variable by t (time) instead of x , and the constant c by P_0 because substitution of $t = 0$ yields $P(0) = P_0$.

We refer to P_0 as the initial population.

Example 3

Suppose that the future population of Stockton City t years after January 1, 2000 is described (in thousands) by the quadratic model

$$P(t) = 110 + 4t + 0.07t^2.$$

- (a) What is the population of Stockton City on January 1, 2007?
- (b) In what month of what calendar year will the population of Stockton City reach 180 thousand?

Quadratic Functions and Modeling

Solution

(a) We only need to substitute $t = 7$ in the population function $P(t)$ and calculate

$$P(7) = 110 + 4(7) + 0.07(7)^2 = 141.43 \text{ thousand.}$$

(b) We need to find the value of t when $P(t) = 180$. That is we need to solve the equation,

$$110 + 4t + 0.07t^2 = 180 \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad (i)$$

Equation (i) is an example of a **quadratic equation** which we can solve in a **variety of ways**.

First, by **graphing** both sides of the equation (figure 3):

$$Y_1 = 110 + 4t + 0.07t^2$$

$$Y_2 = 180$$




figure 3

The line $Y_2 = 180$ and the parabola $Y_1 = 110 + 4x + 0.07x^2$ are shown in the calculator window $-100 \leq x \leq 80$, $-100 \leq y \leq 300$.

To solve equation (i), we find the x -coordinate of the intersection point in the first quadrant. The negative solution of the intersection point in the second quadrant would be in the past. Figure 3 indicates that we have already used the intersection-finding feature to locate the point (14.047, 180).

Hence, $t = 14.047$ years
 $= 14 \text{ years} + (0.047 \times 12) \text{ months}$
 $= 14 \text{ years} + 0.56 \text{ month.}$

Thus Stockton City should reach a population of 180 thousand 14 years, 0.56 month after January 1, 2000.

Hence, sometime during January 2014.

5
Quadratic Functions and Modeling

Alternatively, by using the **Quadratic Formula**:

The quadratic equation $ax^2 + bx + c = 0$, $a \neq 0$
has solutions $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

To use the quadratic formula, we first write equation (i) in the form

$$0.07t^2 + 4t - 70 = 0 \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad (ii)$$

Here, $a = 0.07$, $b = 4$, $c = -70$, giving

$$\begin{aligned} t &= \frac{-4 \pm \sqrt{4^2 - 4(0.07)(-70)}}{2(0.07)} \\ &= \frac{-4 \pm \sqrt{35.6}}{0.14} = 14.046954 \quad \text{or} \quad -71.189811 \end{aligned}$$

The negative solution would be in the past. So, we only accept the positive solution, $t = 14.047$ years = 14 years, 0.56 month.

The Position Function (Model) of a Particle Moving Vertically:

If an object is projected straight upward at time $t = 0$ from a point y_0 feet above ground, with an initial velocity v_0 ft/sec, then its height above ground after t seconds is given by

$$y(t) = -16t^2 + v_0t + y_0.$$

Example 4

A projectile is fired vertically upward from a height of 600 feet above the ground, with an initial velocity of 803 ft/sec.

- (a) Write a quadratic model for its height $y(t)$ in feet above the ground after t seconds.
- (b) During what time interval will the projectile be more than 5000 feet above the ground?
- (c) How long will the projectile be in flight?

Solution

- (a) Given $y_0 = 600$ ft and $v_0 = 803$ ft/sec,

$$y(t) = -16t^2 + 803t + 600.$$

- (b) We need to find the values of t for which the height $y(t) > 5000$ feet.
That is

$$-16t^2 + 803t + 600 > 5000 \quad \cdot \cdot \cdot (i)$$

We solve the inequality (i) by graphing (**figure 4**) both the parabola $Y_1 = -16t^2 + 803t + 600$ and the line $Y_2 = 5000$, in the calculator window $-20 \leq x \leq 70$ and $-2000 \leq y \leq 12000$.




figure 4

The parabola in figure 4 shows the height of the projectile at time t and the horizontal line is the graph of a height of 5000 feet.

The projectile is more than 5000 feet above the ground when the graph of the parabola is above the horizontal line.

Using the intersection-finding feature of the calculator, we find the approximate intersection points to be $t = 6.3$ sec and $t = 43.9$ sec.

Hence, the projectile is above 5000 feet when

$$6.3 \text{ sec} < t < 43.9 \text{ sec}.$$

- (c) The projectile will be in flight until its height $h(t) = 0$. This corresponds to the x -intercept (not the origin) in figure 4.

Using the root or zero feature of the calculator, we obtain

$$t = 50.9 \text{ sec}.$$

EXERCISES

1. The population (in thousands) for Alpha City, t years after January 1, 2004 is modeled by the quadratic function $P(t) = 0.3t^2 + 6t + 80$. In what month of what year does Alpha City's population reach twice its initial (1/1/2004) population?
2. The population (in thousands) for Beta City, t years after January 1, 2005 is modeled by the quadratic function $P(t) = 0.7t^2 + 12t + 200$. How long will it take Beta City's population to reach 350 thousand?
3. The population (in thousands) for Gamma City, t years after January 1, 2002 is modeled by the quadratic function $P(t) = 1.5t^2 + 21t + 300$. How long will it take Gamma City's population to reach 500 thousand?
4. The population (in thousands) for Delta City, t years after January 1, 2003 is modeled by the quadratic function $P(t) = 0.5t^2 + 7t + 90$. In what month of what year does Alpha City's population reach twice its initial (1/1/2003) population?
5. The population (in thousands) for Omega City, t years after January 1, 2002 is modeled by the quadratic function $P(t) = 0.25t^2 + 5t + 100$. In what month of what year does Alpha City's population reach 200 thousand?
6. A ball is thrown straight up, from ground zero, with an initial velocity of 48 feet per second. Find the maximum height attained by the ball and the time it takes for the ball to return to ground zero.
7. From the top of a 48 feet tall building, a ball is thrown straight up with an initial velocity of 32 feet per second. Find the maximum height attained by the ball and the time it takes for the ball to hit the ground.
8. A ball is thrown straight up from the top of a 160 feet tall building with an initial velocity of 48 feet per second. The ball soon falls to the ground at the base of the building. How long does the ball remain in the air?
9. A ball is dropped from the top of a 960 feet tall building. How long does it take the ball to hit the ground?
10. Joshua drops a rock into a well in which the water surface is 300 feet below ground level. How long does it take the rock to hit the water surface?

Quadratic Functions and Modeling

Fitting Quadratic Models to Data:

Find the quadratic model $P(t) = P_0 + bt + at^2$ (with $t = 0$ for the earliest year given in the data)

that best fits the population census data in Problems 11 – 16.

In each case, calculate the average error of this optimal model, and use the model to predict the population in the year 2007.

11. Iowa City, IA

t (years)	1970	1980	1990	2000
P (people)	46,850	50,508	59,735	62,220

12. Arizona

t (years)	2000	2001	2002	2003	2004
P (thous)	5,131	5,320	5,473	5,581	5,744

13. Florida

t (years)	1970	1980	1990	2000	2004
P (thous)	6,789	9,746	12,938	15,982	17,346

14. Georgia

t (years)	1970	1980	1990	2000	2004
P (thous)	4,590	5,463	6,478	8,186	8,825

15. Nevada

t (years)	1970	1980	1990	2000	2004
P (thous)	489	801	1,202	1,998	2,315

16. U.S.

t (years)	1970	1980	1990	2000	2004
P (millions)	203	227	249	281	294

Answers to Exercises

1. February, 2013.
2. 8 years, 143 days.
3. 6 years, 183 days.
4. February, 2011.
5. May, 2014.
6. 36 feet, 3 sec.
7. 64 feet, 3 sec.
8. 5 sec.
9. 7.75 sec.
10. 4.33 sec.
11. $P(t) = 46,234 + 641t - 2.9t^2$; 1377 *people* ; 66,000 *people*.
12. $P(t) = 5,139 + 176t - 6.9t^2$; 14*thous* ; 6,030*thous*.
13. $P(t) = 6,777 + 298t + 0.37t^2$; 44*thous* ; 18,300*thous*.
14. $P(t) = 4,610 + 59t + 1.93t^2$; 59*thous* ; 9,400*thous*.
15. $P(t) = 502 + 14t + 1.16t^2$; 31*thous* ; 2,600*thous*.
16. $P(t) = 204 + 2t + 0.02t^2$; 1.4*thous* ; 300*thous*.

Note: Answers to questions 11-16 are approximate values.

References

1. *Elementary Mathematical Modeling: Functions and Graphs*.
Mary Ellen Davis & C. Henry Edwards. Prentice Hall, 2001.
2. *Intermediate Algebra*.
R. David Gustafson & Peter D. Frisk. Thomson (Brooks/Cole), 2005.
3. www.fairus.org
4. www.gpec.org
5. www.johnson-county.com